

## APPENDIX 1

Relative frequencies (percentages) of the foraminifera in the xenoliths from the Salt Breccia Member of the Wieliczka deposit

Taxon	Foraminiferal biostratigraphy <sup>(*)</sup> and number of sample <sup>(**)</sup>																																															
	IIβ				IIC								IID																																			
	41	45	46	88	95	101	103	30	17	18	1	91	92	93	94	96	97	98	77	99	100	102	63	106	104	105	16	74	76	43	42	71	72	73	47	48	89	90										
BENTHIC	<i>Dendrolyra latissima</i> Grzybowski, 1898																																															
	<i>Reophax varioculus</i> , Luczkowska, 1990																																															
	<i>Spiroplectinella carinata</i> (d'Orbigny, 1846)	0.68	9.46	2.53	0.50	0.67	2.54	0.16	1.01	3.46	1.33	2.34	0.48	0.37	0.62	0.74	0.39	0.85	5.62	3.08	0.26																											
	<i>Pseudotriplasia minuta</i> (Reuss, 1867)										2.28	5.99	7.99	7.37					1.29	3.40	13.07																											
	<i>Karreriella gaudryinoides</i> (Fornasini, 1885)										1.71																																					
	<i>Martinottiella communis</i> (d'Orbigny, 1926)	0.27	3.38	2.02	4.83						18.98	0.26	0.48	0.12																																		
	<i>Textularia gramen</i> d'Orbigny, 1846	0.54		1.01							1.14	0.26	0.24						1.65	9.25																												
	<i>Spirolucina badenensis</i> d'Orbigny, 1846	0.27			0.17						1.68				0.86				0.77																													
	<i>Hauerinidae</i>	0.27	0.68	1.52							2.12	0.67	2.30		0.29	2.28	0.52	0.97	0.50	0.45	0.41	2.02	2.70	0.43	0.61	1.98	1.91	0.39	0.34	0.16	1.35	2.07	2.80	0.16	0.30	2.31	1.56	0.45	0.92	0.55	0.30							
	<i>Nodosaria</i> spp.	5.71	6.76	16.16	9.50						4.38	0.49	2.02	14.41	1.33	8.46	9.69	7.12	0.45	5.19		17.53	5.53	6.69																								
	<i>Frondicula annularis</i> d'Orbigny, 1846	0.14	0.51	0.33															1.21	3.62																												
	<i>Dimorphina variabilis</i> (Neugeboren, 1851)																																															
	<i>Lenticulina ornata</i> (d'Orbigny, 1846)	2.04	6.76	2.53	3.33						0.85	1.18	0.49	1.01	0.29	0.95	0.13	0.48																														
	<i>Sarcenaria arcuata</i> (d'Orbigny, 1846)	0.27																																														
	<i>Marginulina regularis</i> d'Orbigny, 1846										1.01	0.33																																				
	<i>Lagenidae</i>	0.27									0.67	0.42	1.01	0.33	0.29	0.26	0.48	0.12	0.41	0.58	0.30	0.74	1.32	0.26	0.34	0.31	0.37																					
	<i>Guttulina communis</i> d'Orbigny, 1826		0.68								0.51				0.29	0.78	0.48	0.12	0.41	0.58	0.30	0.74	1.32	0.26	0.34	0.31	0.37																					
	<i>Ellipsolagena</i> idae										0.51				0.29																																	
	<i>Glandulina ovula</i> d'Orbigny, 1826										0.17				0.34	0.33																																
	<i>Hoeglundina elegans</i> (d'Orbigny, 1826)																	6.90	3.15	1.50		2.39	0.19		0.61		0.65	4.06																				
	<i>Bolivina</i> spp.	8.02	7.43	14.45	2.67	2.24	5.08	1.85	0.16	1.01	0.29	0.38	0.13	1.94	2.12					0.58	0.61	10.72	0.26	1.69	0.63	1.84		1.31	0.63	0.50	1.16	10.14	0.15	3.30	0.09		6.31											
	<i>Cassidulina laevigata</i> d'Orbigny, 1826	0.14	4.73	7.07	0.33			3.03	0.99		0.29	1.33	1.43	11.86	12.73	5.83	1.04	2.57	5.20	1.70	5.93	3.22	0.29	0.91	8.23	9.42	24.02	0.64	20.52	0.95	27.52	19.21	4.15	0.45	2.41	0.37	0.73	0.45	27.18									
	<i>Globocassidulina crassa</i> (d'Orbigny, 1839)	1.90	11.49	3.54	0.67			5.72	0.33	1.01	3.75	0.38	0.13	3.15	6.24																																	
	<i>Ehrenbergina serrata</i> Reuss, 1850	1.49			3.67																																											
	<i>Bulimina aculeata</i> d'Orbigny, 1826	1.22	0.68	0.51	0.17						0.49	1.01			1.33	2.73	2.91	2.00	0.90	2.28	12.87	0.39	2.98	1.37	1.24	2.98	3.61	6.75	4.53	12.88	7.46	2.85	13.16	0.46	0.15	1.21	2.75	1.29	30.94									
	<i>Bulimina elongata</i> d'Orbigny,																																															