


Fossil karst in the Jurassic of the Kościuszko Mound in Kraków (southern Poland): reply





¹ Polish Geological Institute - National Research Institute. Carpathian Branch, Skrzatów 1, 31-560 Kraków, Poland

Wójcik, A., Garecka, M., Olszewska, B., Wódka., M., 2015. Fossil karst in the Jurassic of the Kościuszko Mound in Kraków (southern Poland): reply. Geological Quarterly, **59** (2): 438–439, doi: 10.7306/gq.1235

The main allegation presented by our opponents (Krajewski and Matyszkiewicz, 2015) is the lack of reference to publications dealing with the Upper Jurassic sediments, especially authored and co-authored by them (Matyszkiewicz, 1996; Krajewski, 2001; Krajewski et al., 2011; Matyszkiewicz et al., 2012; Kochman and Matyszkiewicz, 2013). Our paper did not deal, however, with the development of the Upper Jurassic limestones. The aim of our investigations was to examine the occurrence of the fossil karst at considerable depths within these limestones, and thus the lack of reference to the above-listed literature is not an error. We are simply referring to publications dealing with the problem being the aim of our paper. Certainly, we could describe these limestones in more detail but this lack does not impact the resolution of the problem studied by us, leading to the conclusions of several generations of submarine sedimentation in the karst cavities of the St. Bronisława Hill – unless our opponents think that changes in the limestone structure had a major control on the development of the karst and subsequent filling. We do not find any mention on the occurrence of younger sediments in the Jurassic limestones in any paper quoted by Krajewski and Matyszkiewicz (2015). On the other hand, the earlier papers concerning karst and occurrences of the neptunian dykes in the Kraków area have been referred to (Gradziński, 1962, 1972; Felisiak, 1992; Dżułyński, 1995; Felisiak, 1995; Wieczorek et al., 1995a, b; Kołodziej et al., 2010; Krobicki et al., 2008). Previous studies did not report the marine sedimentation in the caves in Jurassic limestones, as indicated by rudimentary layering both of Miocene limestones and clays with Cretaceous microfauna. We documented a complex and unevenly aged development of the karst system. The reported facts show that the filling of the karst cavities took place during the submarine sedimentation and was not the result of redeposition of the Cretaceous sediments (not to mention Miocene limestones). It does not imply that it cannot occur together with other types filling the karst forms.

Considering the remark on the occurrence of cherts (Krajewski and Matyszkiewicz, 2015: fig. 3) we think that it is not possible to observe their uneven distribution in the borehole, and it can be seen that the cherts which occurred in the cores do not relate probably to the bedding surface because none of them were placed within the interlayer surface. The thickness of the Oxfordian deposits in the Kraków region is not up to 200 m, as we wrote, but more: 230 m according to Matyszkiewicz (1993) and 250 m according to Krajewski et al. (2011). We agree that the title may suggest to some readers that the Kościuszko Mound is built of Jurassic limestones, which was not our intention. We appreciate the exhaustive lecture on ammonite stratigraphy of the Kraków region but the absence of any ammonites in the studied profiles did not allow us for the their reference to the appropriate scheme. It seemed to us that, due to the occurrence of the fault system cutting the discussed region, limestones of different ages can occur at the same depth (as already stressed by Krajewski, 2001). We continued our investigation after the paper had been accepted, and thus our knowledge is more complete now: Rumanolina seiboldi (Lutze) and Protomarssonella jurassica (Mityanina) have been identified in supplementary samples. These are the marker species for Upper Jurassic deposits of the Carpathian Foredeep and Foreland. The former species occurs from the Middle Oxfordian, and the latter is characteristic for the Late Oxfordian. The species have been photographed. In conclusion, the investigated fossil karst forms occur in the upper Oxfordian limestones, and the comments of Krajewski and Matyszkiewicz (2015) do not refer to the main issue of our paper.

Received: March 16, 2015; accepted: April 27, 2015; first published online: May 27, 2015

^{*} Corresponding author, e-mail: antoni.wojcik@pgi.gov.pl

REFERENCES

- Dżułyński, S., 1995. Neptunian dykes of Bonarka a testimony of the Late Cretaceous tectonic movements in the Cracow Upland – discussion (in Polish with English summary). Przegląd Geologiczny, 43: 689.
- Felisiak, I., 1992. Oligocene–early Miocene karst deposits and their importance for recognition of the development of tectonics and relief in the Carpathian Foreland, Kraków region, Southern Poland (in Polish with English summary). Annales Societatis Geologorum Poloniae, 62: 173–207.
- Felisiak, I., 1995. Neptunian dykes of Bonarka a testimony of the Late Cretaceous tectonic movements in the Cracow Upland – discussion (in Polish with English summary). Przegląd Geologiczny, 43: 869–872.
- Gradziński, R., 1962. Origin and development of subterranean karst in the southern part of the Cracow Upland (in Polish with English summary). Rocznik Polskiego Towarzystwa Geologicznego, 32: 429–490.
- **Gradziński, R., 1972**. Przewodnik geologiczny po okolicach Krakowa. Wyd. Geol., Warszawa.
- Kochman, A., Matyszkiewicz, J., 2013. Experimental method for estimation of compaction in the Oxfordian bedded limestones of the southern Kraków–Częstochowa Upland, Southern Poland. Acta Geologica Polonica, 63: 681–696.
- Kołodziej, B., Szulc, J., Machaniec, E., Kędzierski, M., Duda, M., 2010. Injection dykes as evidence of Campanian synsedimentary tectonics on the Kraków Swell, southern Poland. Annales Societatis Geologorum Poloniae, 80: 285–301.
- **Krajewski, M., 2001**. Upper Jurassic chalky limestones in the Zakrzówek Horst, Kraków-Wieluń Upland, Kraków region, south Poland. Annales Societatis Geologorum Poloniae, **71**: 43–52.
- Krajewski, M., Matyszkiewicz, J., 2015. Fossil karst in the Jurassic of the Kościuszko Mound in Kraków (southern Poland): discussion. Geological Quarterly, 59 (2): 433–437.

- Krajewski, M., Matyszkiewicz, J., Król, K., Olszewska, B., 2011.
 Facies of the Upper Jurassic–Lower Cretaceous deposits from the southern part of the Carpathian Foredeep basement in the Kraków–Rzeszów area (southern Poland). Annales Societatis Geologorum Poloniae, 81: 269–290.
- Krobicki, M., Golonka, J., Kucharski, J. 2008. Jurajskie i kredowe dajki neptuniczne w północnej Tetydzie i na obszarze pery-Tetydy reperkusje geotektoniczne (in Polish). Geologia, 34: 185–188.
- Matyszkiewicz, J., 1993. Stratygrafia jura. In: Objaśnienia do Szczegółowej Mapy Geologicznej Polski 1:50 000, arkusz Kraków (973) (ed. J. Rutkowski) (in Polish). Państwowy Instytu Geologiczny, Warszawa.
- Matyszkiewicz, J., 1996. The significance of Saccocoma-calciturbidites for the analysis of the Polish epicontinental Late Jurassic basin: an example from the southern Cracow-Wieluń Upland (Poland). Facies, 34: 23–40.
- Matyszkiewicz, J., Kochman, A., Duś, A., 2012. Influence of local sedimentary conditions on development of microbialites in the Oxfordian carbonate buildups from the southern part of the Kraków–Częstochowa Upland (south Poland). Sedimentary Geology, 263–264: 109–132.
- Wieczorek, J., Dumont, T., Boullin, J., Olszewska, B., 1995a. Neptunian dykes of Bonarka – a testimony of the Late Cretaceous tectonic movements in the Cracow Upland – reply (in Polish with English summary). Przegląd Geologiczny, 43: 690–692.
- Wieczorek, J., Dumont, T., Bouillin, J., Olszewska, B., 1995b. Neptunian dykes of Bonarka – a testimony of the Late Cretaceous tectonic movements in the Cracow Upland – reply (in Polish with English summary). Przegląd Geologiczny, 43: 872-875.