

APPENDIX 6

Chemical EMP composition of iron-nickel metal in host rock, clasts and unequilibrated inclusions in the Pułtusk breccia

[wt.%]	H3.8 inclusions						H4/H5 host rock			Clasts H5 and H6		
P	b.d.	b.d.	b.d.	b.d.	0.01	b.d.	b.d.	b.d.	b.d.	b.d.	b.d.	b.d.
Fe	95.61	75.93	64.12	63.87	67.59	78.56	71.03	93.06	50.37	93.99	66.34	48.16
Ni	4.44	22.28	35.64	35.87	32.83	20.62	28.78	6.41	49.14	4.58	33.31	50.79
Co	0.46	0.50	0.04	0.03	0.06	1.04	0.14	0.45	0.04	0.55	0.21	0.07
Cu	b.d.	0.04	0.14	0.11	0.09	0.04	0.16	0.03	0.36	b.d.	b.d.	0.34
S	0.01	b.d.	b.d.	b.d.	0.02	0.02	b.d.	b.d.	0.01	b.d.	b.d.	b.d.
Zn	b.d.	b.d.	b.d.	b.d.	b.d.	b.d.	b.d.	0.01	b.d.	0.03	b.d.	b.d.
Total	100.52	98.75	99.94	99.88	100.60	100.28	100.11	99.96	99.92	99.15	99.86	99.36

b.d. – below detection limit