

^{224}Ra and the $^{224}\text{Ra}/^{228}\text{Ra}$ activity ratio in selected mineral waters from the Polish Carpathians

Chau NGUYEN DINH¹, Lucyna RAJCHEL^{1,*}, Hao DUONG VAN² and Jakub NOWAK³

¹ AGH University of Science and Technology, Faculty of Geology, Geophysics and Environmental Protection, al. A. Mickiewicza 30, 30-059 Kraków, Poland

² Hanoi University of Mining and Geology, Faculty of Oil and Gas, Vietnam

³ AGH University of Science and Technology, Faculty of Physics and Applied Computer Science, al. A. Mickiewicza 30, 30-059 Kraków, Poland



Nguyen Dinh, Ch., Rajchel, L., Duong Van, H., Nowak, J., 2017. ^{224}Ra and the $^{224}\text{Ra}/^{228}\text{Ra}$ activity ratio in selected mineral waters from the Polish Carpathians. *Geological Quarterly*, **61** (4): 771–778, doi: 10.7306/gq.1365

Mineral and therapeutic waters widely occurring in the Polish Carpathians contain a vast amount of biogenic elements. However, radioactive elements like radium isotopes are also found in these waters. This paper presents the first results of the activity concentration measurements of ^{224}Ra in selected mineral waters of the Polish Carpathians. Additionally, the content of ^{226}Ra and ^{228}Ra , and the activity ratios of $^{224}\text{Ra}/^{228}\text{Ra}$, $^{224}\text{Ra}/^{226}\text{Ra}$ and $^{226}\text{Ra}/^{228}\text{Ra}$ in the waters were determined. The work was based on a method for the simultaneous determination of ^{224}Ra , ^{226}Ra and ^{228}Ra in water samples using the liquid scintillation spectrometer. The results show that the activity concentrations of ^{224}Ra , ^{226}Ra and ^{228}Ra in the mineral waters are comparable and vary from ca. 220 mBq/L to above 1500 mBq/L. The activity ratios of $^{224}\text{Ra}/^{228}\text{Ra}$, $^{224}\text{Ra}/^{226}\text{Ra}$ and $^{226}\text{Ra}/^{228}\text{Ra}$ are variable within the ranges of 0.68 to 1.48, 0.65 to 1.48, and 0.78 to 2.05, respectively. The committed effective dose resulting from the intake of ^{224}Ra through the water consumption is far lower than that from ^{226}Ra and ^{228}Ra .

Key words: ^{224}Ra , radium, mineral waters, effective dose, LSC method.

INTRODUCTION

The determination of radium isotopes in water samples is very important from the hydrological and radiological point of view. The relation between the radium isotopes' concentration in groundwater and the lithology of the host aquifers were reported by many scientists (e.g., King et al., 1982; Sturchio et al., 1992; Martin and Akber, 1999; Lucivjansky, 1999; Ruberu et al., 2005; Grundl and Cape, 2006; Walencik-Lata et al., 2016). The thermodynamic properties and applications of the radium isotopes for interaction between groundwater and aquifer rock were studied by Davidson and Dickson (1986), Langmuir and Melchior (1985) and Szabo et al. (2012). The contribution of ^{226}Ra and ^{228}Ra to annual committed effective dose resulting from the consumption of mineral and drinking waters amounts to >70% (Bayés et al., 1996; Manjón et al., 1996; Godoy et al.,

2001; Waller and Steininger, 2007; Waller et al., 2008; Nguyen et al., 2012).

According to the Council Directive (Euratom, 2013), drinking water is safe from the radiological point of view if with the consumption of two litres per day, the annual committed dose rate originating from the presence of the radioactive elements in this water is <0.1 mSv. In general, in many countries (for example in Poland) there are natural artesian water intakes serving as direct sources of drinking water for population. In that case, the ^{224}Ra should be taken into account in the estimation of committed effective doses.

The Carpathians are the richest region of mineral and therapeutic waters in Poland, where numerous spas and bottling plants are located. A significant number of mineral waters occurring in this region are available for consumers directly from boreholes or springs. The main aim of the work was the determination of ^{224}Ra content in selected mineral waters from the Polish Carpathians. For the first time, the results of the activity concentration of ^{224}Ra in the Polish Carpathian mineral waters are presented. Additionally, the content of other radium isotopes (^{226}Ra and ^{228}Ra) and the activity ratios of $^{224}\text{Ra}/^{228}\text{Ra}$, $^{224}\text{Ra}/^{226}\text{Ra}$ and $^{226}\text{Ra}/^{228}\text{Ra}$ were determined. In the paper, some geochemical and radiological aspects of the presence of three radium isotopes (^{224}Ra , ^{226}Ra and ^{228}Ra) and their activity ratios in studied mineral waters are also discussed.

* Corresponding author, e-mail: lucynar@agh.edu.pl




Fig. 1. Map of mineral waters sampling sites

SAMPLING AND ANALYSIS METHOD

STUDY REGION AND SAMPLING

The water samples were collected from several mineral water intakes in the Polish Outer Carpathians (Fig. 1). This region is composed chiefly of Cretaceous and Miocene sandstones and mudstone of flysch origin, in which carbonate and siliceous rocks sporadically occur. The formations are strongly folded and divided into different units. Due to the very complicated geological structure, the Carpathians are regarded as a region of diverse hydrogeological conditions (e.g., Rajchel, 2012; Rajchel and Czop, 2012). This region hosts many valuable mineral water resources used for balneotherapy and by mineral water plants. The samples were collected from some selected resources of therapeutic bicarbonated waters, waters containing carbon dioxide, and chloride waters from the Magura Unit (Zuber-1 and Zuber-2 boreholes, and Słotwinka spring in Krynica-Zdrój; Krakus, Warzelnia and Rabka IG 2 boreholes in Rabka-Zdrój) and the Silesian Unit (Elin 7 and Emma boreholes in Iwonicz-Zdrój; Klimkówka 27 borehole in Klimkówka; Lubatówka 12 borehole in Lubatówka; Table 1). The waters are diagenetic and mixed with the infiltration water to different degrees (Rajchel and Czop, 2012).

In the Magura Unit (Inoceranian Bed; Fig. 1) the average uranium and thorium concentrations are 2.06 and 8.92 ppm, but in the Silesian Unit (Godula, Istebna, Ciężkowice and Krosno beds; Oszczypko et al., 2006) they vary from 1.07 to 1.98 and 4.03 to 10.05, respectively (Plewa and Plewa, 1992).

The water samples were placed into 2-litre plastic bottles. Before sampling the bottles were rinsed by HCl 0.1 M and distilled water. To eliminate the remaining liquid, the bottle was rinsed several times with the water intended for analysis at the

water intake. The sampling time was recorded and the samples were transported to the laboratory as soon as possible.

METHODOLOGY

At the laboratory, the water samples were weighed and the radium isotopes were co-precipitated from the water sample together with barium as a sulphate compound. The precipitate was placed into the glass vial of volume 22 mL, mixed with 12 mL of the gel scintillation cocktail, and measured by the *Guardian alpha/beta Wallac Scintillation Counter* (LSC method). For every sample, the measurement was performed daily for two hours. The measurement process lasted until the radioactive equilibrium between ^{226}Ra and ^{222}Rn has been established. The applied chemical preparation and measurement procedures were described in detail by Nguyen et al. (1997).

After radium precipitation, only radium isotopes (^{226}Ra , ^{228}Ra and ^{224}Ra) occur in the obtained sample. ^{226}Ra belongs to the uranium series with ^{238}U as a primary element, ^{228}Ra and ^{224}Ra are the members of thorium series with ^{232}Th as a first isotope. At the radioactive equilibrium occurring in the ^{226}Ra group, there are four alpha decay isotopes (^{226}Ra , ^{222}Rn , ^{218}Po , ^{214}Po) and two beta decay isotopes (^{218}Pb , ^{214}Bi). In the ^{228}Ra group, there are only two beta decay isotopes (^{228}Ra , ^{228}Ac). The ^{224}Ra group emits four alpha and two beta particles. Four alpha particles are related to ^{224}Ra , ^{220}Rn , ^{216}Po (every isotope emits one particle per decay), and ^{212}Bi and ^{212}Po with emission of 0.337 per decay and 0.663 per decay, respectively. Two beta particles are emitted from ^{212}Pb (one per decay), ^{212}Bi (0.663 per decay) and ^{208}Tl (0.337 per decay; Brown et al. 1986). Since the half-life ($T_{1/2}$) of ^{226}Ra is 1620 y, and the longest daughter in this group is ^{222}Rn with $T_{1/2}$ equal to 3.825 d, so the secular equilibrium in the group is established after nearly 500 h

Table 1

Selected data on boreholes, springs and medicinal waters

Locality/type and name of water intake	Depth [m]	Depth of the water intake [m]	Water type	GPS coordinates
Krynica-Zdrój Zuberb-1 borehole	810.0	696.2–810.0	HCO ₃ -Na+CO ₂	591.2 m a.s.l. 49°24'50"N 20°57'48"E
Krynica-Zdrój Zuber-2 borehole	948.5	586.8–670.0	HCO ₃ -Na-Mg+CO ₂	576.2 m a.s.l. 49°24'53"N 20°57'44"E
Krynica-Zdrój Słotwinka spring	–	–	HCO ₃ -Mg-Na-Ca+CO ₂ +Fe	615.0 m a.s.l. 49°25'59"N 20°57'25"E
Rabka-Zdrój Warzelnia borehole	50.2	41.0–45.7	Cl-Na+Fe+I	494.99 m a.s.l. 49°36'36"N 19°58'02"E
Rabka-Zdrój Rabka IG 2 borehole	1215.0	1185–1215	Cl-Na+I+T	563.84 m a.s.l. 49°36'57"N 19°58'30"E
Rabka-Zdrój Krakus borehole	19.0	19.0	Cl-Na+I	495.5 m a.s.l. 49°36'35"N 19°57'58"E
Klimkówka Klimkówka 27 borehole	1255.0	417.0–481.0	HCO ₃ -Cl-Na+CO ₂ +I	462.8 m a.s.l. 49°33'26"N 21°47'32"E
Iwonicz-Zdrój Emma borehole	283.7	102.8–263.0	Cl-HCO ₃ -Na+CO ₂ +I	435.20 m a.s.l. 49°33'38"N 21°46'58"E
Iwonicz-Zdrój Elin 7 borehole	1030.0	85.0–238.0	Cl-HCO ₃ -Na+CO ₂ +I	434.80 m a.s.l. 49°33'39" N 21°45'02"E
Lubatówka Lubatówka 12 borehole	1151.5	625.0–958.0	Cl-HCO ₃ -Na+CO ₂ +I+T	350.7 m a.s.l. 49°33'59"N 21°47'32"E

from the moment of radium precipitation. In the ²²⁸Ra group, the activity of ²²⁸Ac can reach the ²²⁸Ra activity in about 20 h after the precipitation. With reference to the lab time (one month), the ²²⁸Th can be regarded as a "stable" isotope due to its *T*_{1/2} equal to 1.9 y. The *T*_{1/2} of ²²⁴Ra is 3.62 d and the longest lived progeny is ²¹²Pb with the *T*_{1/2} = 10.6 h, so the transient equilibrium in the group is established after 53 h. Since that moment, the total alpha and beta activities of the ²²⁴Ra group decrease in accordance with decay law. After 500 h, the total activity of the ²²⁴Ra group is below 2% of the activity of this isotope at the precipitation time. Therefore, the contribution of the ²²⁴Ra group to the total alpha and beta count rates after 500 h elapsed from the radium precipitation can be neglected, and the ²²⁶Ra and ²²⁸Ra can be estimated using the alpha and beta count rates measured after 500 h from the moment of radium precipitation. For every studied sample the dependence of the net alpha and beta intensities (the intensities of the background sample were subtracted from the adequate intensities of the sample) on the time elapsed from the precipitating moment was used to determine the ²²⁴Ra, ²²⁶Ra and ²²⁸Ra contents.

The ²²⁶Ra concentration in a water sample is estimated by the formula:

$$^{226}\text{Ra} = \frac{I_{500}}{F_{500}^{Ra\ 226} \cdot 60} \cdot V \quad [1]$$

where: *I*₅₀₀ – net count rate (cpm) of the sample measured in the alpha channel after 500 h from the moment of radium precipitation; *F*₅₀₀^{Ra 226} – factor corresponding to relative alpha activity of the ²²⁶Ra group at 500 h elapsed from the sample preparation; this factor can be obtained from the Bateman equation or empirically using the sample prepared from the ²²⁶Ra standard solution; – the detection efficiency of the liquid scintillation detector for alpha particle and can be estimated using the sample of ²²⁶Ra standard solution or arbitrarily taken to be equal to 100% (Horrocks, 1974); *V* – water sample volume; the 60 is the number of seconds in one minute.

The ²²⁸Ra concentration in the water sample is determined using the beta count rate. The total beta count rate at 500 h elapsed time consists of both beta count rates from the ²²⁸Ra and ²²⁶Ra groups. The count rate of the beta particles emitted from the ²²⁶Ra group is calculated by the following formula:

$$I_{500}^{Ra\ 226} = I_{500} \cdot C \quad [2]$$

where: *C* = $\frac{I_{500}^{st}}{I_{500}^{st}}$; *I*₅₀₀st and *I*₅₀₀st is the count rates for the ²²⁶Ra standard sample after 500 h from the moment of radium precipitation measured in the beta and alpha channels, respectively.

The count rate from the ²²⁸Ra group in the sample is equal to:

$$I_{500}^{Ra\ 228} = I_{500}^{total} - I_{500}^{Ra\ 226} \quad [3]$$

where: I_{500}^{total} – the count rate measured in the beta channel for the sample.

The ^{228}Ra concentration in water sample is estimated by the formula:

$$^{228}\text{Ra} = \frac{I_{500}^{\text{Ra } 228}}{F_{500}^{\text{Ra } 228} \cdot 60 \cdot V} \quad [4]$$

where: $F_{500}^{\text{Ra } 228}$ – the factor corresponding to the relative beta activity of the ^{228}Ra group after 500 h from the moment of radium precipitation, ϵ – the efficiency of the detection of beta particles emitted from the ^{228}Ra group. The other symbols denote the same as in Formula [1].

The count rates measured daily in the alpha channel consist of the count rate originating from both the ^{226}Ra and ^{224}Ra groups, so the count rate measured at the t -moment from the ^{224}Ra content in the prepared water sample is determined as follows:

$$I^{\text{Ra } 224}(t) = I^{total}(t) - I^{\text{Ra } 226}(t) \quad [5]$$

where: $I^{total}(t)$, $I^{\text{Ra } 226}(t)$ are the total alpha count rate and the alpha count rate originating from the ^{226}Ra group, respectively; t – the elapsed time from the moment of radium precipitation.

The $I^{\text{Ra } 226}(t)$ can be calculated by the formula:

$$I^{\text{Ra } 226}(t) = I_0^{\text{Ra } 226} \cdot F^{\text{Ra } 226}(t) \quad [6]$$

where: $I_0^{\text{Ra } 226}$ – the activity concentration determined by Formula [1], $F^{\text{Ra } 226}(t)$ is the factor corresponding to the relative alpha activity for the ^{226}Ra group at the time (t) elapsed from the sample.

$F^{\text{Ra } 226}(t)$ can be calculated using the Bateman equation or determined empirically using the sample prepared from the ^{226}Ra standard solution. The spectra and spectra of the prepared ^{228}Ra standard sample measured at 50 and 265 h elapsed from the radium precipitation are presented in Figure 2 and Figure 3, respectively. The figures show that the alpha activity decreased in agreement with the ^{224}Ra decay and the beta activity decreased clearly more slowly in comparison with the alpha one.




Fig. 2. Alpha and beta spectra of the ^{228}Ra standard solution measured by Wallac Winspectral 1414 alpha and beta liquid scintillation counter at 50 h elapsed from the radium precipitation




Fig. 3. Alpha and beta spectra of the ^{228}Ra standard solution measured by Wallac Winspectral 1414 alpha and beta liquid scintillation counter at 265 h elapsed from the radium precipitation




Fig. 4. Total alpha count rates and its components of ^{226}Ra and ^{224}Ra groups for the water sample collected from the Zuber-2 intake (for the fitting curve value of $\lambda = 0.0081 \text{ h}^{-1}$ corresponding to $T_{1/2}$ equal to 3.57 days)

This fact is related to both the constant beta activity of the ^{228}Ra group and the decreasing beta activity of the ^{224}Ra group. The measured total alpha count rate as a function of the elapsed time and its separated components originating from the ^{226}Ra and ^{224}Ra groups for the sample collected from the Zuber-2 water intake is shown in Figure 4 as an example. In order to estimate the ^{224}Ra content, the curve (according to the decay form $-ae^{-bt}$) presenting the relation between the component of the alpha count rate of the ^{224}Ra group and time elapsed from the precipitating moment (after 53 h) is fitted. The content of ^{224}Ra is calculated using the decay law of the ^{224}Ra group and $I_0^{\text{Ra } 224}$ which is an ordinate of the intersection point of the above-mentioned fitted curve on the y-coordinate.

$$^{224}\text{Ra} = \frac{I_0^{\text{Ra } 224} \cdot e^{-\lambda_{\text{Ra } 224} \cdot 53}}{60 \cdot F_{53}^{\text{Ra } 224} \cdot V} \cdot e^{-\lambda_{\text{Ra } 224} \cdot t} \quad [7]$$

$$= \frac{I_0^{\text{Ra } 224} \cdot e^{-\lambda_{\text{Ra } 224} \cdot t}}{60 \cdot F_{53}^{\text{Ra } 224} \cdot V}$$

where: λ_{Ra-224} [h^{-1}] – the fitted value of the ²²⁴Ra decay constant, and F_{53}^{Ra-224} the factor corresponding to the relative alpha activity of the ²²⁴Ra group at 53 hours elapsed from the precipitating water sample, and can be estimated using the Bateman equation for the ²²⁴Ra group or determined empirically using the sample prepared from the ²³²Th standard solution, where there is radioactive equilibrium between ²²⁸Ra and ²²⁴Ra. t [h] – the time elapsed from the water sampling moment to the precipitation point.

In accordance with the definition described by Currie (2008) and propagation law, the limit of detection was calculated based on the average uncertainty of the count rates in the alpha and beta channels for the background sample, and is equal to 11 mBq/dm³ for ²²⁴Ra, 7 mBq/dm³ for ²²⁶Ra, and 30 mBq/dm³ for ²²⁸Ra.

RESULTS AND DISCUSSION

The activity concentrations of ²²⁴Ra, ²²⁶Ra and ²²⁸Ra in the studied waters vary from 206 to 1116 mBq/L, from 213 to 1545 mBq/L, and from 186 to 752 mBq/L, respectively (Table 2). Figures 5, 6 and 7 show the relationships between ²²⁶Ra and ²²⁴Ra, between ²²⁸Ra and ²²⁴Ra, and between ²²⁸Ra and ²²⁶Ra, respectively. All points (excluding the water from Lubatówka 12) in the figures are scattered between line 1:1. There are strong linear correlations between the studied radium isotopes. The Pearson correlation coefficient for the relations between ²²⁶Ra and ²²⁴Ra, between ²²⁸Ra and ²²⁴Ra, and between ²²⁸Ra and ²²⁶Ra is equal to 0.94, 0.88 and 0.85, respectively. The three analysed isotopes are variants of the same chemical element (radium); thus, their chemical properties and behaviour in groundwater are the same.

The activity ratios of ²²⁴Ra/²²⁸Ra and ²²⁶Ra/²²⁸Ra range in the intervals of 0.68–1.48 and 0.78–2.05, respectively. The av-

erage value of ²²⁴Ra/²²⁸Ra activity ratio is 1.09, and, if we exclude the water from Lubatówka 12, this ratio decreases to 1.04. The observed phenomena indicate that the activity concentration of ²²⁴Ra in groundwater is generally slightly higher than that of ²²⁸Ra, which is related to the recoil effect. In the thorium decay chain, ²²⁴Ra is followed by two alpha decays, while ²²⁸Ra is followed only by one alpha decay. Therefore, ²²⁴Ra shows higher probability of migration due to direct transfer of the atom across the solid/liquid phase boundary or lattice destruction by recoil effect than ²²⁸Ra. It is the same process which causes uranium isotopes disequilibrium (the activity ratio of ²³⁴U/²³⁸U is generally >1) in groundwater (Osmond and Cowart, 1976; Nguyen et al., 2011).

In such mineral waters, the chemical processes dominate the recoil effect and radium isotopes migrate mainly from host rocks to water due to leaching processes. The lower content of ²²⁴Ra than ²²⁸Ra can be associated with the fact that thorium (²²⁸Th is progeny of ²²⁸Ra and parent of ²²⁴Ra) forms low-soluble and highly adsorbed compounds in contrast to radium (Langmuir, 1997; Nguyen et al., 2011).

The observed average activity ratio values (²²⁶Ra/²²⁸Ra) for all studied waters are 1.2, being typical for highly mineralised groundwater in the Polish Carpathians (Nguyen et al., 2016). Generally, the activity concentrations of ²²⁶Ra, ²²⁸Ra and their activity ratio (²²⁶Ra/²²⁸Ra) in groundwater are related to the adequate content and activity ratio of their primordial isotopes (²³⁸U, ²³²Th, ²³⁸U/²³²Th) in the host aquifers (Asikainen and Kahlos, 1979; King et al., 1982; Vesterbacka et al., 2006; Nguyen et al., 2012). For six waters from the Magura Unit and three waters from the Silesian Unit, the average activity concentration ratios (²²⁶Ra/²²⁸Ra) are 1.04 and 1.47, respectively. The values correspond, to a certain degree, to the ratios of the average uranium and thorium concentrations in the Magura and Silesian units. The highest activity concentrations of the radium isotopes and their activity ratios (²²⁶Ra/²²⁸Ra and ²²⁴Ra/²²⁸Ra) were observed

Table 2

The activity concentrations of ²²⁴Ra, ²²⁶Ra and ²²⁸Ra and their activity ratios in the studied Carpathian mineral waters

No.	Locality/ water intake	TDS [g/L]	Activity concentration [mBq/L]			Activity concentration ratio		
			²²⁴ Ra	²²⁶ Ra	²²⁸ Ra	²²⁴ Ra/ ²²⁸ Ra	²²⁴ Ra/ ²²⁶ Ra	²²⁶ Ra/ ²²⁸ Ra
1	Krynica-Zdrój Zuber-1	23.2	428 38	568 24	530 38	0.81	0.75	1.07
2	Krynica-Zdrój Zuber-2	20.5	460 32	447 23	312 34	1.47	1.03	1.43
3	Krynica-Zdrój Słotwinka	3.52	420 17	284 19	362 30	1.16	1.48	0.78
4	Rabka-Zdrój Warzelnia	20.6	216 17	314 19	312 25	0.70	0.69	1.01
5	Rabka-Zdrój Rabka IG 2	23.5	722 45	619 25	598 39	1.21	1.17	1.04
6	Rabka-Zdrój Krakus	23.8	331 23	442 30	485 41	0.68	0.75	0.91
7	Klimkówka Klimkówka 27	12.2	218 34	333 15	215 25	1.01	0.65	1.55
8	Iwonicz-Zdrój Emma	5.92	206 13	213 8	203 18	1.01	0.97	1.05
9	Iwonicz-Zdrój Elin 7	5.80	245 23	231 17	186 19	1.31	1.06	1.24
10	Lubatówka Lubatówka 12	20.5	1116 71	1545 88	752 61	1.48	0.72	2.05




Fig. 5. ²²⁶Ra concentration vs. ²²⁴Ra concentration in the water samples




Fig. 7. ²²⁶Ra concentration vs. ²²⁸Ra concentration in the water samples




Fig. 6. ²²⁸Ra concentration vs. ²²⁴Ra concentration in water samples




Fig. 8. Activity concentrations of radium isotopes (²²⁴Ra, ²²⁶Ra and ²²⁸Ra) vs. TDS in the water samples

in the water from Lubatówka 12. This fact can be related to the local geology and hydrogeological conditions in this area.

The TDS in the investigated groundwater varies from 3.5 to 23.8 g/L. The relations between activity concentrations of three radium isotopes (²²⁴Ra, ²²⁶Ra and ²²⁸Ra) and TDS are presented in Figure 8. There is no clear relation between the radium content in investigated waters and TDS, or it is difficult to notice because of the small amount of data.

Based on the obtained activity concentrations of the radium isotopes, the annual committed effective doses (*E*) resulting from consumption of two litres of the mineral waters a day for adults were calculated using the following equation:

$$E = V \sum_{i=1}^N A_i DCF_i \quad [8]$$

where: *V* stands for consumption of water (2L per day), *A_i* – the activity concentration of the given radionuclide in water expressed in

Bq/L, and *DCF_i* stands for the DCFs (Dose Conversion Factor) for ingestion by humans belonging to the given age group (adults in this case). The DCF is equal to the dose in Sv resulting from the ingestion of 1Bq of the given radionuclide (ICRP, 1995; WHO, 2008).

The calculated committed effective dose from ²²⁶Ra and ²²⁸Ra changes in the range from 44 to 316 μSv/year and from 94 to 379 μSv/year, respectively. The data show that the dose from the ²²⁴Ra varies from 7.5 to 53 μSv/year and its contribution to the total annual committed effective dose is <8% (Table 3). Thus, the intake of ²²⁴Ra with consumed water does not pose a significant radiological risk to human health in comparison with other two radium isotopes (²²⁶Ra and ²²⁸Ra).

CONCLUSIONS

The activity concentrations of ²²⁴Ra in the studied mineral waters are comparable to that of ²²⁶Ra and ²²⁸Ra. There is a

Table 3

The annual committed effective doses for adults resulting from intake of the waters, calculated for three radium isotopes (^{224}Ra , ^{226}Ra and ^{228}Ra)

No.	Town/ water intake	Annual committed effective dose [μSv]			Contribution of ^{224}Ra to the total effective dose [%]
		^{224}Ra	^{226}Ra	^{228}Ra	
1	Krynica-Zdrój Zuber-1	15	116	267	3.8
2	Krynica-Zdrój Zuber-2	20	86	302	7.1
3	Krynica-Zdrój Slotwinka	15	58	182	6.0
4	Rabka-Zdrój Warzelnia	8.4	64	157	3.7
5	Rabka-Zdrój Rabka IG 2	34	127	301	7.4
6	Rabka-Zdrój Krakus	12	90	244	3.6
7	Klimkówka Klimkówka 27	11	68	108	5.9
8	Iwonicz-Zdrój Emma	7.5	44	102	4.9
9	Iwonicz-Zdrój Elin 7	10	47	94	6.7
10	Lubatówka Lubatówka 12	53	316	379	7.1

strong linear correlation between ^{224}Ra and the other two radium isotopes.

The average value of the observed activity ratio ($^{224}\text{Ra}/^{228}\text{Ra}$) is slightly above 1 (1.04). This is related to the recoil effect; the same process which causes the disequilibrium of uranium isotopes in groundwater. The activity ratio of $^{224}\text{Ra}/^{228}\text{Ra}$ lower than 1 can be related to the fact that thorium forms low-soluble and highly adsorbed compounds in an aqueous environment. The activity ratios of $^{226}\text{Ra}/^{228}\text{Ra}$ in the Magura and Silesian units are related to the average concentrations of uranium and thorium in water-bearing formations occurring in the units.

The intake of ^{224}Ra with consumed water does not pose a significant radiological risk to human health in comparison with ^{226}Ra and ^{228}Ra .

The LSC method is clearly useful for the simultaneous determination of ^{224}Ra , ^{226}Ra and ^{228}Ra in water samples.

Acknowledgements. The authors would like to express their thanks to the anonymous reviewers for their valuable remarks. This work was supported by the AGH University of Science and Technology projects Nos.: 11.11.140.645, 11.11.140.862 and 15.11.220.717.

REFERENCES

- Asikainen, M., Kahlos, H., 1979.** Anomalously high concentrations of uranium, radium and radon in water from drilled well in the Helsinki region. *Geochimica et Cosmochimica Acta*, **44**: 1681–1686.
- Bayés, J.C., Gómez, E., Garcias, F., Casas, M., Cerdá, V., 1996.** Radium determination in mineral waters. *Applied Radiation and Isotopes*, **47**: 849–853.
- Brown, E., Firestone, R., Shirley, V., 1986.** Table of Radioactive Isotopes. John Wiley & Sons, New York, Chichester, Brisbane, Toronto, Singapore.
- Currie, L.A., 2008.** Detection and quantification capabilities in nuclear analytical measurement. In: *Analysis of Environmental Radionuclides* (ed. S.P. Povinec): 49–136. Elsevier.
- Davidson, M.R., Dickson, B.L., 1986.** A porous flow model for steady-state transport of radium in ground water. *Water Resources Research*, **22**: 34–44.
- Euratom, 2013.** Council Directive 2013/51/Euratom of 22 October 2013 laying down requirements for the protection of the health of the general public with regard to radioactive substances in water intended for human consumption.
- Godoy, J.M., da Amaral, E.C., Luiza, M., Godoy, D.P., 2001.** Natural radionuclides in Brazilian mineral water and consequent doses to the population. *Journal of Environmental Radioactivity*, **53**: 175–182.
- Grundl, T., Cape, M., 2006.** Geochemical factors controlling radium activity in sandstone aquifer. *Ground Water*, **44**: 518–527.
- Horrocks, D.L., 1974.** Application of Liquid Scintillation Counting. Academic Press, New York.
- ICRP, 1995.** Age-Dependent Doses to Members of the Public From Intake of Radionuclides – Part 5 Compilation of Ingestion and Inhalation Dose Coefficients. International Commission on Radiological Protection, Publication 72, Ann. ICRP 26 (1).
- King, P., Michel, J., Moore, W., 1982.** Ground water geochemistry of ^{228}Ra , ^{226}Ra and ^{222}Rn . *Geochimica et Cosmochimica Acta*, **46**: 1173–1182.
- Langmuir, D., 1997.** Aqueous Environmental Chemistry. Prentice Hall. Upper Saddle River, New Jersey.

- Langmuir, D., Melchior, D., 1985.** The geochemistry of Ca, Sr, Ba and Ra sulfates in some deep brines from Palo Duro Basin, Texas. *Geochimica et Cosmochimica Acta*, **49**: 2423–2432.
- Lucivjansky, L., 1999.** Natural radioactivity of water in Slovakia. *Slovak Geological Magazine*, **5**: 53–61.
- Manjón, G., Vioque, I., Moreno, H., García-Tenorio, R., García-León, M., 1996.** Determination of ^{226}Ra and ^{224}Ra in drinking waters by liquid scintillation counting. *Applied Radiation and Isotopes*, **48**: 535–540.
- Martin, P., Akber, R.A., 1999.** Radium isotopes as indicators of adsorption-desorption interactions and barite formation in groundwater. *Journal of Environmental Radioactivity*, **46**: 271–286.
- Nguyen, D.C., Niewodniczański J., Dorda J., Ochoński A., Chruściel E., Tomza I., 1997.** Determination of radium isotopes in mine waters through α - and β -activities measured by liquid scintillation spectrometry. *Journal of Radioanalytical & Nuclear Chemistry*, **222**: 69–74.
- Nguyen, D.C., Duliński, M., Jodłowski, P., Nowak, J., Różański, K., Śleziak, M., Wachniew, P., 2011.** Natural radioactivity in groundwater - a review. *Isotopes in Environmental and Health Studies*, **47**: 415–437.
- Nguyen, D.C., Rajchel, L., Nowak, J., Jodłowski, P., 2012.** Radium isotopes in the Polish Outer Carpathian mineral waters of various chemical composition. *Journal of Environmental Radioactivity*, **112**: 38–44.
- Nguyen, D.C., Kopeć, M., Nowak J., 2016.** Factors controlling ^{226}Ra , ^{228}Ra and their activity ratio in groundwater – an application in Polish Carpathian mineral waters. *Geology, Geophysics & Environment*, **42**: 337–351.
- Osmond, J.K., Cowart, J.B., 1976.** The theory and uses of natural uranium isotopic variations in hydrology. *Atomic Energy Review*, **14**: 621–679.
- Oszczypko, N., Uchman, A., Malata, E., 2006.** Rozwój paleotektoniczny basenów Karpat zewnętrznych i pienińskiego pasa skałkowego (in Polish). Instytut Nauk Geologicznych Uniwersytetu Jagiellońskiego, Kraków.
- Plewa, M., Plewa S., 1992.** *Petrofizyka* (in Polish). Wyd. Geol., Warszawa.
- Rajchel, L., 2012.** Carbonated waters and waters containing carbon dioxide of the Polish Carpathians (in Polish with English summary). Wydawnictwo AGH, Kraków.
- Rajchel, L., Czop, M., 2012.** Hydrogeochemical modelling of chloride mineral water from Rabka spa (Carpathians Mountains, Poland). *Geological Quarterly*, **56** (4): 681–690.
- Ruberu, S.R., Liu, Y.G., Perera, S.K., 2005.** Occurrence of ^{224}Ra , ^{226}Ra , ^{228}Ra , gross alpha, and uranium in California groundwater. *Health Physics*, **89**: 667–678.
- Sturchio, N.C., Bohlke, J.K., Markun, F.J., 1992.** Radium isotope geochemistry of thermal water, Yellowstone National Park, Wyoming, USA. *Geochimica et Cosmochimica Acta*, **57**: 1203–1214.
- Szabo, Z., DePaul, V.T., Fischer, J.M., Kraemer, T.F., Jacobsen, E., 2012.** Occurrence and geochemistry of radium in water from principal drinking-water aquifer system of the United States. *Applied Geochemistry*, **37**: 729–752.
- Vesterbacka, P., Turtiainen, T., Heinävaara, S., Arvela, H., 2006.** Activity concentrations of ^{226}Ra and ^{228}Ra in drilled well water in Finland. *Radiation Protection Dosimetry*, **121**: 406–421.
- Walencik-Łata, A., Kozłowska, B., Dorda, J., Przylibski, T.A., 2016.** The detailed analysis of natural radionuclides dissolved in spa water of the Kłodzko Valley, Sudety Mountains, Poland. *Science of the Total Environment*, **569–570**: 1174–1184.
- Waller, G., Steininger, G., 2007.** Radium isotopes and ^{222}Rn in Austrian drinking waters. *Journal of Radioanalytical and Nuclear Chemistry*, **274**: 511–516.
- Waller, G., Wagner, R., Katzlberger, Ch., 2008.** Natural radionuclides in Austrian mineral water and their sequential measurement by fast methods. *Journal of Environmental Radioactivity*, **99**: 1090–1094.
- WHO, 2008.** Guidelines for Drinking-Water Quality. World Health Organization, Geneva.