

### APPENDIX 3

#### Indices and proxies of foraminifera calculated for samples from GSS IV (index of sample as in Appendix 1)

Sample	Percentage of benthic	Number of benthic taxa	Percentage of benthic ecogroups *)				Central Paratethys biozone
			epibenthic	inbenthic	organic-rich	opportunistic	
70	97.1	12	6.8	92.8	53.3	29.4	CPN 8
69	46.4	10	2.9	96.7	5.5	89.9	
68	44.8	17	8.0	91.0	18.9	51.5	
67	27.9	6	3.8	95.1	33.9	45.4	
66	89.6	10	20.9	79.1	27.9	36.3	
65	12.1	10	4.2	95.8	9.7	72.9	
54	96.2	9	6.0	80.9	49.4	10.4	
53	17.4	2	0.0	100.0	0.0	97.4	
40	41.0	17	8.5	79.3	17.5	62.7	
39	98.0	9	11.8	76.2	47.3	19.0	
38	51.9	12	4.0	95.1	1.8	92.7	
36	3.6	7	3.7	92.6	81.5	7.4	
35	39.9	11	3.2	94.7	59.1	34.8	
34	79.3	19	2.7	95.1	65.1	11.7	
33	51.6	12	7.0	88.8	33.0	14.7	
29	98.3	15	6.4	93.1	54.8	8.9	
27	9.7	6	0.0	88.3	15.0	70.0	
26	54.4	16	8.6	89.4	40.7	40.0	
21	90.6	12	7.4	92.6	41.5	48.0	
20	48.4	15	11.0	71.3	33.6	26.5	
19	97.2	15	4.9	94.6	73.9	21.0	
15	4.7	2	0.0	100.0	3.2	96.8	
56	29.9	31	19.3	74.0	26.4	14.9	CPN 7
37	28.6	18	22.1	69.5	49.5	11.6	

\* – details in the text