

New ornithischian dinosaur footprints in the Jurassic of Poland

Gerard GIERLI SKI, El bieta GA DZICKA, Grzegorz NIED WIEDZKI and Grzegorz PIE KOWSKI

Gierli ski G., Ga dzicka E., Nied wiedzki G. and Pie kowski G. (2001) — New ornithischian dinosaur footprints in the Jurassic of Poland. *Geol. Quart.*, 45 (2): 205–210. Warszawa.

New stratigraphic horizons with the ornithischian footprints have been found in central Poland. Basal thyreophoran ichnites are reported from the middle Hettangian Skłoby Formation of Gromadzice. They support the previous conclusion that small and medium-sized Early Jurassic ornithischians preferred a coastal habitat with low-rise vegetation. Another find, an ornithopod footprint, came from the early Kimmeridgian Głowaczów Formation of O arów quarry. The track was discovered within the shallow-marine succession of calcareous deposits, which contributes to the interpretation of synsedimentary emersion of that area during the early Kimmeridgian.

Gerard Gierli ski, El bieta Ga dzicka and Grzegorz Pie kowski, Polish Geological Institute, Rakowiecka 4, PL-00-975 Warszawa, Poland, e-mail: gpie@pgi.waw.pl; Grzegorz Nied wiedzki, Piotrowice 91/5, PL 23-107 Strzy ewice, Poland, e-mail: GrzegorzNiedziedzki@poczta.net-line.pl (received: February 14, 2001; accepted: March 8, 2001).

Key words: Poland, Hettangian, Kimmeridgian, Ornithischia, dinosaurs, footprints.

INTRODUCTION

The largest ornithischian track assemblage in Poland is known from the late Hettangian coastal barrier-lagoon deposits of Gliniany Las (Gierli ski and Potemska, 1987; Gierli ski, 1991, 1999; Gierli ski and Pie kowski, 1999). Less numerous specimens of ornithischian footprints have been also found in the similar coastal palaeoenvironments of the Hettangian of Zapniów (Pie kowski and Gierli ski, 1987) and in so-called lower Gromadzice outcrop (Gierli ski and Pie kowski, 1999). Those previous finds led to the supposition that small and medium-sized Jurassic ornithischians preferred a coastal (barrier-lagoon and delta plain) habitat with low and dense vegetation rather than the inland forest habitat invaded by high browsing adult sauropods (Gierli ski and Pie kowski, 1999). The discovery of ornithischian tracks in the upper Gromadzice outcrop supports such a viewpoint.

The second find reported herein, the ornithopod footprint from the Upper Jurassic of O arów, indicates a temporary emergence of that area during the early Kimmeridgian times and correspond with the coastal environment supposed to be a part of the ornithischian habitats.

THYREOPHORAN FOOTPRINT FROM GROMADZICE

The upper Gromadzice outcrop (Fig. 1B) of the deltaic middle Hettangian deposits of the Skłoby Formation (Pie kowski, 1991) hitherto revealed only unusually small tracks of diminutive or juvenile sauropods (Gierli ski, 1997). Recently two ornithischian footprints have been found in this horizon.

The outcrop reveals complete section of a delta sequence (Fig. 1C). This particular deltaic environment was characterised by flat prodelta slope, intermediate wave energy, considerable littoral drift and intermediate discharge of fluvial sediments. It resulted in the “composite sequence”, in which the delta cycle is divided into two parts separated by the barrier (delta-front) facies formed in the wave-breaking zone. The “inner delta facies” are represented by deposits formed in stagnant lagoons (interdistributary bays) and marshes, covered by quickly prograding distributary (fluvial) facies (Pie kowski, 1991). The footprints were preserved on the bottom surface of the gray sandstone representing distributary channel facies. The distributary channel facies covers the clayey coal bed with dense plant roots beneath (palaeosoil). It points that the animal entered an area with relatively high water-table level and




Fig. 1. **A** — Location of the studied area

B — Geological sketch of the northeastern slope of the Holy Cross Mountains (HCM)

P — Permian, T — Triassic, J₁ — Lower Jurassic, J₂ — Middle Jurassic, J_o — Oxfordian, J_k — Kimmeridgian, K — Cretaceous

C — Sedimentological section of the Hettangian deposits in the upper Gromadzice outcrop (after Pie kowski, 1991)

1 — parallel bedding, 2 — large-scale cut-and-fill trough cross-bedding, 3 — small- and medium-scale trough cross-bedding, 4 — hummocky cross stratification, 5 — wave ripples, 6 — flaser bedding, 7 — wavy bedding, 8 — lenticular bedding, 9 — parallel lamination in mudstones, 10 — plant roots, 11 — coal and coal-bearing mudstone; **lithologic scale**: m. — mudstone, st. — siltstone, sd. — sandstone, c — presence of mud clasts; **facies**: o.s. — offshore, s.f. — shoreface, p.d. — prodelta, f.d.-b. — delta front-barrier, b.b. — backbarrier, L-i.b.-m. — lagoon-interdistributary bay-marsh system, d.ch. — distributary channel

D — Simplified lithostratigraphic section of the Kimmeridgian deposits in O arów quarry (after Gutowski, 1998)

12 — biogenic and intraclastic grainstones and packstones, 13 — clays and marls, 14 — oyster shell-beds, 15 — micritic limestones, 16 — oolites, 17 — chalky limestones, 18 — ammonite (d — *Divisum*, h — *Hypselocyclum*), 19 — dinosaur footprints; **lithologic units**: Oopl — O arów oolite and platy limestone, Ool — O arów oyster lumachelle

low-rise vegetation. Erosion at the bottom of the distributary channel was insignificant due to the weak currents, which increased the preservation potential of the tracks.

Both ornithischian footprints are preserved as natural casts in a sandstone slab (MUZ PIG OS-221/35). They may belong to the same trackway or to two parallel trackways. The ichnites show a typical anomoepodid pattern, a functionally tridactyl pes with relatively short and highly divaricated digits. However, they have exceeded the typical *Anomoepus* size, reaching the length of 20.7 cm and showing only two phalangeal pads on digit III. Their morphology fits almost perfectly between the more gracile tracks of *Anomoepus Hitchcock*, 1848 and the more robust *Moyenisauropus karaszewskii* Gierli ski, 1991. Such a morphological pattern has been described from the Lower Jurassic of Lesotho as *Moyenisauropus natator* Ellenberger, 1974 and it has been also reported from the Lower Jurassic, from the Kayenta-Navajo transition zone of Utah (Lockley *et al.*, 1998a).

Some authors consider Ellenberger's *Moyenisauropus* as junior synonym of *Anomoepus* (Olsen and Galton, 1984; Thulborn, 1994). Gierli ski (1991) agreed that most of *Moyenisauropus* tracks are indeed *Anomoepus*, except for *M. natator* and possibly *M. levicauda*. Recently, Lockley and Meyer (2000) contended that all seven ichnospecies of *Moyenisauropus* should be transferred to *Anomoepus*, while the Polish form of *M. karaszewskii* should be classed as a completely different, new ichnogenus. However, the present authors share the opinion of Gierli ski and Pie kowski (1999). *Moyenisauropus natator* is morphologically intermediate between *Anomoepus* and *M. karaszewskii*. Even if *M. karaszewskii* would fall into a different ichnogenus, the ichnogenus *Moyenisauropus sensu* Gierli ski (1991) would be valid to comprise such an intermediate form.

The discussed sample of *M. natator* from the upper Gromadzice outcrop (Fig. 2A) shows digit divarication as follows: I–II = 86°, II–III = 36°, III–IV = 27°, II–IV = 63°. Digits II and III are subequal in length, whereas digit IV is the longest. Their length ratios measured according to the method of Olsen *et al.* (1998) equal III/II = 1.10 and III/IV = 0.62, thus fitting closely those of the foot of *Scelidosaurus harrisonii* Owen, 1861 (see Gierli ski and Pie kowski, 1999, tab. 1).

ORNITHOPOD FOOTPRINT FROM O ARÓW

The second ichnite reported herein, the specimen MUZ PIG 1663.II.2, came from the Late Jurassic limestones of O arów quarry (Fig. 1B). The tridactyl footprint is a well preserved natural cast of the right pes impression (Fig. 2B). This print demonstrates morphology clearly different from the Early Jurassic *Anomoepus-Moyenisauropus* morphotypes. The imprint is nearly as long as wide (18 cm long and 19 cm wide), very symmetrical with almost the same angle between the axes of digit II and III (40°) as between digit III and IV (41°). Very similar ornithopod footprints from the Kimmeridgian Morrison Formation of Utah have been named *Dinehichnus* by Lockley *et al.* (1998b). However, the type material from Utah shows the "heel" pad impression, a distinctive metatarsal-phalangeal pad

of the fourth digit, more circular and discrete than the Polish specimen. If the shallow posteromedial mark was produced there by a hallux, it might be another character differing it from the typical *Dinehichnus* tracks.

The digit length ratios (III/II = 1.35 and III/IV = 0.75) of the Polish ichnite resemble those of supposed dryosaur footprint from the Morrison Formation of Como Bluff in Wyoming (III/II = 1.32 and III/IV = 0.78) reported by Bakker (1996). However, the presence of probable hallux imprint in the O arów specimen speaks against its dryosaur origin, suggesting rather camptosaurid affinity of its trackmaker.

Highly intriguing is the palaeoenvironmental background of this find. Ornithischian dinosaurs were not demonstrably associated with aquatic environments and Bakker (1996) even concluded that the compact-footed herbivores such as camptosaurus and dryosaurus preferred rather areas with well-drained soil than swampy terrain. However, the sediment that has preserved (casted) the discussed footprint has been evidently deposited in marine environment.

During the Late Jurassic the northeastern margin of the Holy Cross Mts. (HCM) as well as the surrounding regions of the Central Poland were predominantly covered by shallow epicontinental seas. The nearest lands situated to the south-east (the Ukrainian shield) and to the north (the Baltic shield) lied some hundreds kilometres away from the HCM (Ziegler, 1990). During the middle Oxfordian an extensive carbonate ramp developed on the stable basement of the East European Craton, probably close to the Ukrainian shield (Kutek, 1994). It prograded to the north-west and (together with smaller ones that grew on the local elevations) was a source of carbonates in the Polish Basin. Thus from the middle Oxfordian until the early Kimmeridgian sedimentation was dominated by carbonates on the substantial part of this area. Different types of calcareous rocks with biogenic, intraclastic, micritic and oolitic limestones, coral and sponge-algal bioherms, gaizes and marls build today a huge series of deposits, reaching approximately 600 metres thickness in the neighbourhood of O arów quarry. Lithological diversity and a presence of stratal discontinuities suggest changes in depositional regime during this time and allow to distinguish sedimentary sequences of various significance. Depositional sequences can also be recognised on geophysical logs (e.g. gamma rays) of boreholes drilled in this region like B kowa IG1 and Ciepiałów IG1 (Niemczycka, 1974, 1975). The find of ornithischian dinosaur footprint suggests a regional surface of depositional discontinuity.

The surface into which the dinosaur track was printed has not been found. Thus, there is no possibility of checking up if it shows any signs of erosional structures, karstification or pedogenic modification. The natural cast of the ornithopod footprint found in the O arów quarry came from the shallow-marine succession of mostly calcareous deposits. A fifty centimetres thick layer of pale micritic limestone was qualified by M. Poł ska from the Polish Geological Institute (PGI) as a mudstone/wackestone. Besides carbonates it contains quartz (about 2%) and mica grains, ferrum oxides and hydroxides, aggregates of some opaque minerals, lidites and plant remains. The microfossil analysis, kindly done by J. Smole (from PGI), revealed content of benthic foraminifers shells as well as very abundant fragments of echinoderms. Among the foraminifers




Fig. 2. **A** — *Moyensauropus natator* Ellenberger, 1974 from the Early Jurassic Skłoby Formation of Gromadzice, MUZ PIG OS-221/35A; **B** — cf. *Dinehichmus* sp. from the Late Jurassic Głowaczów Formation of O arów, MUZ PIG 1663.II.2

dominant ones are these of the genus *Spirillina* (*S. tenuissima* Gumbel) and *Trocholina*. The whole sequence, about fifty eight metres thick, contains also cross-bedded grainstones and intercalations of oolites (Fig. 1D). It was distinguished by Gutowski (1998) as the "O arw oolite and platy limestone" (Oopl) in his informal lithostratigraphic scheme. This author designated also an early Kimmeridgian age for that sequence corresponding to the ammonite *Hypselocyclum* Zone. The faunal content does not allow to accurately date the boundary between the Oxfordian and Kimmeridgian in the mentioned section. Within the superimposed sequence of lumachelle limestone (Oll), the *Divisum* Zone of the lower Kimmeridgian was recognised (Gutowski, 1998). Earlier in the northwestern part of the Lublin Upland and adjacent area of the HCM Niemczycka (1976) distinguished some formal lithostratigraphic units. A shallow-water succession containing calcareous and dolomitic oolites, lumachelles, intraclastic and oncolithic limestone as well as storm induced deposits has been named the Gowaczw Formation. The early Kimmeridgian age have also been indicated for it. D browska (1983) emphasised an extremely shallow-water (nearshore) environment for the early Kimmeridgian sedimentation in the vicinity of H a that is situated some forty kilometres north-westwards to O arw. Moreover, she postulated a temporary emersion in this part of the sedimentary basin. In the lower Kimmeridgian sedimentary succession, she observed abundant shellbeds, wave related structures and increased input of terrigenous material to the basin resulted in deposition of sand beds and conglomerates.

It is also noteworthy that Liszkowski (1972) found well preserved and abundant plant remains in the Upper Jurassic of this area. Besides branches and seeds, they included delicate leaves, pieces of flowers and cones, excluding long transport.

The fossils were attributed to some groups of the Gymnospermyta; the dominance of representatives of the Coniferales pointed to an arid climate. These fossil plants occurred in transgressive shallow-marine succession of carbonate deposits containing basal conglomerate, oolites and platy limestones with lenticular-shape intercalations of oolites. Lithological features of deposits indicate near-shore sedimentation and give evidences for synsedimentary emersion. The author suggested the late Oxfordian age for this sequence, but no fossils of stratigraphic importance have been found there. The succession studied is very similar to these of the lower Kimmeridgian.

Of course, further and more detailed investigations, especially in the terms of sequence stratigraphy, should be carried on in the Upper Jurassic sections of HCM. However, we can conclude, that sedimentary succession of the lower Kimmeridgian in the northeastern margin of the HCM gives evidences of temporary emergence of the area. The dinosaur footprint reveal evidence of a regional sea level fall. It had to be pronounced enough to result in connection of this area with distant land(s). According to Haq *et al.* (1988), the considerable fall of sea level caused by eustasy in the early Kimmeridgian is dated for 144 Ma. This event is correlated with the lower part of boreal ammonite *Cymodoce* Zone. The find of the ornithopod footprint in the lower Kimmeridgian of HCM contributes to the interpretation of sedimentary periodicity and a succession of palaeoenvironments in this region.

Abbreviations of cited repositories: MUZ PIG — Geological Museum of the Polish Geological Institute, Warsaw, Poland; MUZ PIG OS — Holy Cross Mts. Branch of the Polish Geological Institute, Kielce, Poland.

REFERENCE

- BAKKER R. T. (1996) — The real Jurassic Park: dinosaur and habitats at Como Bluff, Wyoming. In: The Continental Jurassic (ed. M. Morales). Mus. Northern Arizona Bull., **60**: 34–49.
- D BROWSKA Z. (1983) — Jura okolic H y. In: Paleontologia i stratygrafia jury i kredy okolic H y. Materiały VII Krajowej Konferencji Paleontologw. H a, 7–9 X 1983: 14–24.
- ELLENBERGER P. (1974) — Contribution a la classification des Pistes de Vertebres du Trias: les types du Stormberg d'Afrique du Sud (II). Palaeovertebrata, Memoire extraordinaire: 1–201.
- GIERLI SKI G. (1991) — New dinosaur ichnotaxa from the Early Jurassic of the Holy Cross Mountains, Poland. Palaeogeogr. Palaeoclimat. Palaeoecol., **85** (1–2): 137–148.
- GIERLI SKI G. (1997) — Sauropod tracks in the Early Jurassic of Poland. Acta Palaeont. Pol., **42** (4): 533–538.
- GIERLI SKI G. (1999) — Tracks of a large thyreophoran dinosaur from the Early Jurassic of Poland. Acta Palaeont. Pol., **44** (2): 231–234.
- GIERLI SKI G. and PIE KOWSKI G. (1999) — Dinosaur track assemblages from the Hettangian of Poland. Geol. Quart., **43** (3): 329–346.
- GIERLI SKI G. and POTEMSKA A. (1987) — Lower Jurassic dinosaur footprints from Gliniany Las, northern slope of the Holy Cross Mountains, Poland. N. Jb. Geol. Palont. Abh., **175** (1): 107–120.
- GUTOWSKI J. (1998) — Oxfordian and Kimmeridgian of the northeastern margin of the Holy Cross Mountains, Central Poland. Geol. Quart., **42** (1): 59–72.
- HAQ B. U., HARDENBOL J. and VAIL P. R. (1988) — Mesozoic and Cenozoic chronostratigraphy and eustatic cycles. In: Sea Level Changes: an Integrated Approach (eds. C. K. VILUS *et al.*). Spec. Publs. Soc. Econ. Paleont. Miner., **42**: 71–108.
- HITCHCOCK E. (1848) — An attempt to discriminate and describe the animals that made the fossil footmarks of the United States, and especially New England. Mem. Am. Acad. Arts. Sc., **3** (2): 129–256.
- KUTEK J. (1994) — Jurassic events in south-eastern cratonic Poland. Acta Geol. Pol., **44** (3–4): 165–221.
- LISZKOWSKI J. (1972) — First Upper Jurassic paleofloristic locality in Poland. Prz. Geol., **20** (8–9): 388–393.
- LOCKLEY M. G., HUNT A. P., MEYER C. A., RAINFORTH E. C. and SCHULTZ R. J. (1998a) — A survey of fossil footprint sites at Glen Canyon National Recreation Area (Western USA): a case study in documentation of trace fossil resources at a National Reserve. Ichnos, **5**: 177–211.
- LOCKLEY M. G., SANTOS V. F., MEYER C. A. and HUNT A. (1998b) — A new dinosaur tracksite in the Morrison Formation, Boundary Butte, southeastern Utah. Modern Geol., **23**: 317–330.

- LOCKLEY M. G. and MEYER C. A. (2000) — Dinosaur Tracks and Other Fossil Footprints of Europe. Columbia University Press. New York.
- NIEMCZYCKA T. (1974) — Jura. In: Ciepeliów IG 1 (ed. T. Niemczycka). Prof. Gł b. Otw. Wiert. Inst. Geol., **20**: 27–39.
- NIEMCZYCKA T. (1975) — Jura górna. In: B kowa IG 1 (ed. T. Niemczycka). Prof. Gł b. Otw. Wiert. Inst. Geol., **26**: 23–34.
- NIEMCZYCKA T. (1976) — Lithostratigraphy of Upper Jurassic deposits in the Lublin–Radom area (in Polish with English summary). Acta Geol. Pol., **26** (4): 569–602.
- OLSEN P. E. and GALTON P. M. (1984) — A review of the reptile and amphibian assemblages from the Stormberg of southern Africa with special emphasis on the age of the Stormberg. Palaeont. Afr., **25**: 87–110.
- OLSEN P. E., SMITH J. B. and McDONALD N. G. (1998) — The material of the type species of the classic theropod footprint genera *Eubrontes*, *Anchisauripus* and *Grallator* (Early Jurassic, Hartford and Deerfield basins, Connecticut and Massachusetts, U.S.A.). J. Vertebrate Paleont., **18**: 586–601.
- OWEN R. (1861) — A monograph of the fossil Reptilia of the Lias Formations. Part I: *Scelidosaurus harrisonii*. Palaeont. Soc. Monogr., **13**: 1–14.
- PIE KOWSKI G. (1991) — Eustatically-controlled sedimentation in the Hettangian-Sinemurian (Early Jurassic) of Poland and Sweden. Sedimentology, **38** (3): 503–518
- PIE KOWSKI G. and GIERLI SKI G. (1987) — New finds of dinosaur footprints in Liassic of the Holy Cross Mountains and its palaeoenvironmental background. Prz. Geol., **35** (4): 199–205.
- THULBORN R. A. (1994) — Ornithopod dinosaur tracks from the Lower Jurassic of Queensland. Alcheringa, **18**: 247–258.
- ZIEGLER P. (1990) — Geological Atlas of Western and Central Europe. Shell Intern. Petr. Maatsch. B.V. Hague.